

T 0508 HILL LAB (44 555 22) +64 7 858 2000 E mail@hill-labs.co.nz W www.hill-laboratories.com

## **Certificate of Analysis**

Page 1 of 3

SPv1

Client: Revital Fertilisers

Contact: D Gibson

> C/- Revital Fertilisers PO Box 8045 New Plymouth 4342

Lab No: 2244700 **Date Received:** 20-Sep-2019 27-Sep-2019 **Date Reported:** 


**Quote No:** 98937 **Order No:** 26124 **Client Reference: Drilling Mud Submitted By:** D Gibson

Sample Type: Sludge						
\$	Sample Name:	Turongi 8 12-Sep-2019 12:00 pm				
	Lab Number:	2244700.1				
Individual Tests					I	I
Dry Matter	g/100g as rcvd	76	_	-	_	-
Total Recoverable Barium	mg/kg dry wt	2,900	-	-	-	-
Total Recoverable Potassium	mg/kg dry wt	3,800	-	-	-	-
Total Recoverable Sodium	mg/kg dry wt	810	-	-	-	-
Chloride*	mg/kg dry wt	10,500	-	-	-	-
pH	pH Units	10.68	-	-	-	-
Total Nitrogen*	g/100g dry wt	0.09	-	-	-	-
Heavy metals, screen As,Cd,C						
Total Recoverable Arsenic	mg/kg dry wt	14	_	-	_	-
Total Recoverable Cadmium	mg/kg dry wt	< 0.10	_	-	-	-
Total Recoverable Chromium	mg/kg dry wt	30	-	-	-	-
Total Recoverable Copper	mg/kg dry wt	40	-	-	-	-
Total Recoverable Lead	mg/kg dry wt	92	-	-	-	-
Total Recoverable Mercury	mg/kg dry wt	< 0.10	-	-	-	-
Total Recoverable Nickel	mg/kg dry wt	18	-	-	-	-
Total Recoverable Zinc	mg/kg dry wt	75	-	-	-	-
BTEX in Solids by Headspace						
Benzene	mg/kg dry wt	< 0.06	-	-	-	-
Toluene	mg/kg dry wt	< 0.06	-	-	-	-
Ethylbenzene	mg/kg dry wt	< 0.06	-	-	-	-
m&p-Xylene	mg/kg dry wt	< 0.12	-	-	-	-
o-Xylene	mg/kg dry wt	< 0.06	-	-	-	-
Polycyclic Aromatic Hydrocarb	ons Screening in S	olids				1
1-Methylnaphthalene	mg/kg dry wt	0.77	-	-	-	-
2-Methylnaphthalene	mg/kg dry wt	1.12	-	-	-	-
Acenaphthylene	mg/kg dry wt	< 0.14	-	-	-	-
Acenaphthene	mg/kg dry wt	< 0.14	-	-	-	-
Anthracene	mg/kg dry wt	< 0.14	-	-	-	-
Benzo[a]anthracene	mg/kg dry wt	< 0.14	-	-	-	-
Benzo[a]pyrene (BAP)	mg/kg dry wt	< 0.14	-	-	-	-
Benzo[a]pyrene Potency Equivalency Factor (PEF) NES	mg/kg dry wt	< 0.4	-	-	-	-
Benzo[a]pyrene Toxic Equivalence (TEF)	mg/kg dry wt	< 0.4	-	-	-	-
Benzo[b]fluoranthene + Benzo[fluoranthene	j] mg/kg dry wt	< 0.14	-	-	-	-
Benzo[e]pyrene	mg/kg dry wt	< 0.14	-	-	-	-
Benzo[g,h,i]perylene	mg/kg dry wt	< 0.14	-	-	-	-



Sample Type: Sludge						
	Sample Name:	Turongi 8 12-Sep-2019 12:00 pm				
	Lab Number:	2244700.1				
Polycyclic Aromatic Hydrocarbons Screening in Solids						
Benzo[k]fluoranthene	mg/kg dry wt	< 0.14	-	-	-	-
Chrysene	mg/kg dry wt	< 0.14	-	-	-	-
Dibenzo[a,h]anthracene	mg/kg dry wt	< 0.14	-	-	-	-
Fluoranthene	mg/kg dry wt	< 0.14	-	-	-	-
Fluorene	mg/kg dry wt	< 0.14	-	-	-	-
Indeno(1,2,3-c,d)pyrene	mg/kg dry wt	< 0.14	-	-	-	-
Naphthalene	mg/kg dry wt	0.9	-	-	-	-
Perylene	mg/kg dry wt	< 0.14	-	-	-	-
Phenanthrene	mg/kg dry wt	0.21	-	-	-	-
Pyrene	mg/kg dry wt	< 0.14	-	-	-	-
Total Petroleum Hydrocarbons in Solids						
C7 - C9	mg/kg dry wt	131	-	-	-	-
C10 - C14	mg/kg dry wt	27,000	-	-	-	-
C15 - C36	mg/kg dry wt	46,000	-	-	-	-
Total hydrocarbons (C7 - C36	6) mg/kg dry wt	73,000	-	-	-	-





## **Summary of Methods**

The following table(s) gives a brief description of the methods used to conduct the analyses for this job. The detection limits given below are those attainable in a relatively clean matrix. Detection limits may be higher for individual samples should insufficient sample be available, or if the matrix requires that dilutions be performed during analysis.

Unless otherwise indicated, analyses were performed at Hill Laboratories, 28 Duke Street, Frankton, Hamilton 3204.

Sample Type: Sludge					
Test	Method Description	Default Detection Limit	Sample No		
Environmental Solids Sample Drying*	Air dried at 35°C Used for sample preparation. May contain a residual moisture content of 2-5%.	-	1		
Environmental Solids Sample Preparation	Air dried at 35°C and sieved, <2mm fraction. Used for sample preparation. May contain a residual moisture content of 2-5%.	-	1		
Heavy metals, screen As,Cd,Cr,Cu,Ni,Pb,Zn,Hg	Dried sample, <2mm fraction. Nitric/Hydrochloric acid digestion, ICP-MS, screen level.	0.10 - 4 mg/kg dry wt	1		
Dry Matter (Env)	Dried at 103°C for 4-22hr (removes 3-5% more water than air dry), gravimetry. (Free water removed before analysis, non-soil objects such as sticks, leaves, grass and stones also removed). US EPA 3550.	0.10 g/100g as rcvd	1		
esICextn*	(1:5) ratio of sample (g):0.02M potassium dihydrogen orthophosphate extractant (mL), analysis by Ion Chromatography. In House.	-	1		
Total Recoverable digestion	Nitric / hydrochloric acid digestion. US EPA 200.2.	-	1		

Sample Type: Sludge			
Test	Method Description	Default Detection Limit	Sample No
Total Recoverable Barium	Dried sample, sieved as specified (if required).  Nitric/Hydrochloric acid digestion, ICP-MS, screen level. US EPA 200.2.	0.4 mg/kg dry wt	1
Total Recoverable Potassium	Dried sample, sieved as specified (if required).  Nitric/Hydrochloric acid digestion, ICP-MS, screen level. US EPA 200.2.	100 mg/kg dry wt	1
Total Recoverable Sodium	Dried sample, sieved as specified (if required).  Nitric/Hydrochloric acid digestion, ICP-MS, screen level. US EPA 200.2.	40 mg/kg dry wt	1
Chloride*	Ion Chromatography determination of a potassium phosphate extract of an environmental solid.	3 mg/kg dry wt	1
рН	pH meter. APHA 4500-H+ B 23 <sup>rd</sup> ed. 2017.	0.10 pH Units	1
Total Nitrogen*	Catalytic Combustion (900°C, O2), separation, Thermal Conductivity Detector [Elementar Analyser].	0.05 g/100g dry wt	1

These samples were collected by yourselves (or your agent) and analysed as received at the laboratory.

Samples are held at the laboratory after reporting for a length of time depending on the preservation used and the stability of the analytes being tested. Once the storage period is completed the samples are discarded unless otherwise advised by the client.

This certificate of analysis must not be reproduced, except in full, without the written consent of the signatory.

Ara Heron BSc (Tech)

Client Services Manager - Environmental